

Applied experiences focused on citrus nutrition and water management for mitigating HLB

Experiencias aplicadas en la nutrición
de cítricos y manejo de agua para
mitigar del HLB

Davie Kadyampakeni

University of Florida

Citrus Research and Education Center

Lake Alfred, FL33850.

2º CIL, Guadalajara, Jalisco, Mexico 11 October, 2019



Outline

- Status of HLB in FL: What we know now
- Irrigation studies for managing HLB: Examples
- Nutrition studies for managing HLB: Highlights
- Summary
- Acknowledgements

Current Status of HLB

- Citrus accounts for \$10 billion in economic activity
- Pre-HLB 240 million boxes (10 billion tons)
- Current 80 million boxes (3.3 billion tons), about 67% reduction in production
- Production costs up to \$2100 per acre due to HLB
- Significant reduction in production area
- Declined tree performance, root loss and significant defoliation

Irrigation strategies for managing HLB


- Preventative measures: HLB negative (healthy trees)
 - Frequent irrigation (daily or multiple times a day) e.g. Citrus Under Cover Production System
 - Regulated deficit irrigation
 - Partial root zone drying
 - Plus Asian psyllid control
- Curative management of HLB positive trees (asymptomatic trees)
 - Daily irrigation plus Asian psyllid control
 - Managing pH to optimum levels for nutrient availability
 - Improved nutrition programs via fertigation
- Remediation/Management of HLB affected trees (symptomatic trees)
 - Daily irrigation plus Asian psyllid control
 - Managing pH to optimum levels for nutrient availability
 - Fertigation practices

Irrigation strategies for managing HLB

Field studies on irrigation conducted in:

- Irrigation studies at 3 sites: Ave Maria, Avon Park, Arcadia (2013-2014)
 - ➔ Comparison of Daily, IFAS (recommended by UF) and Intermediate Irrigation Schedules based on Florida Automated Weather Network (FAWN) evapotranspiration
- Advanced Citrus Production Systems (ACPS) studies:
Two Sites: Immokalee at UF/IFAS, SWFREC, and Lake Alfred (2008 to 2011)
 - ➔ Comparison of drip and modified microsprinkler irrigation with grower practices
- Greenhouse studies conducted at Immokalee, SWFREC (2014-2015)
 - ➔ Comparison of HLB vs non-HLB affected citrus

Irrigation studies



- Daily > Intermediate > IFAS irrigation scheduling
- Daily irrigation could help in managing HLB affected trees, reduce tree water stress

Water use of HLB affected trees in south west and central Florida



Irrigation studies

Total available water (%) in southwest and central Florida

Irrigation treatment	Soil depth (cm)	Commercial site		
		Arcadia	Avon Park	Immokalee
Daily				
	0-15	68.9dc	80.7b	68.1bc
	15-30	72.2c	58.7c	75.3b
	30-45	98.2a	87.8a	97.9a
Intermediate				
	0-15	52.2fg	56.3cd	64.5c
	15-30	58.9ef	61.4c	46.6d
	30-45	98.8a	74.3b	42.3d
IFAS				
	0-15	48.1g	49.7d	46.6d
	15-30	80.9b	50.4d	32.1e
	30-45	62.3de	61.9c	69.3bc

- Increasing TAW with depth, greater uptake in the top 6 inches.
- Greater TAW in in top 15 cm than lower 15-45 cm for Daily than Intermediate and IFAS irrigation schedule.

Irrigation studies in central and southwest Florida



Moisture contents and significant relationships with sapflow



Water use of HLB affected trees in southwest Florida under greenhouse conditions

Month -year	ET _o (mm d ⁻¹)	ET _c (mm d ⁻¹)		ET _c diff. (%) [‡]
		Hamlin-Non HLB	Hamlin-HLB	
Jan-Jun-14	3.57	2.97	2.23	23.73
Jul-Dec-14	4.42	4.16	2.63	34.82
Jan-Jun-2015	3.38	4.08	2.83	29.82
Jun-Oct-15	3.73	4.94	3.18	35.20
Overall Average	3.79	4.00a**	2.69b**	30.75
		Valencia-Non HLB	Valencia-HLB	
Jan-Jun-14	3.57	2.83	2.22	22.28
Jul-Dec-14	4.42	3.97	2.83	28.85
Jan-Jun-2015	3.38	3.85	2.69	30.98
Jun-Oct-15	3.73	4.79	3.56	26.42
Overall Average	3.79	3.82a**	2.80b**	26.99**

- 22 to 35% greater water use for Non-HLB affected trees
- Inter-season and annual variability in water use
- Comparable water use between varieties

Citrus Crop Coefficients between HLB and Non-HLB affected Citrus Trees

- Patterns of K_c similar for HLB affected and non-affected trees
- Non-affected tree K_c similar to those found to field trees prior to greening
- Infected trees consistently with lower K_c
- 35.2% in 2014 and 20.8% in 2015




Crop coefficient (K_c) for HLB affected trees in southwest Florida under greenhouse conditions

Nutrition studies for managing HLB: Highlights


- Advanced Citrus Production Systems (ACPS) studies:
 - ➔ Two Sites: Immokalee at UF/IFAS, Immokalee, and Lake Alfred (2008 to 2011)
 - ➔ Comparison of drip and modified microsprinkler fertigation systems with Conventional grower practices
 - ➔ Two ACPS systems: drip (DOHS) and microsprinkler (RM, MOHS), and conventional microsprinkler practice (CMP)

ACPS Nutrition Studies



Leaf NPK concentration

Leaf NPK conc. were in optimum or high range.



Nutrient efficiency (m³/kg N)

CMP needs more fertilizer and water per ha at 1 to 2.5 years than ACPS (Schumann et al. 2010)


ACPS Nutrition Studies (2)

N and P accumulation on Immokalee sand

Fertigation method	CMP	Drip	RM	CMP	Drip	RM
Tissue	N (kg ha ⁻¹)			P (kg ha ⁻¹)		
Leaves	24.00	49.78	37.10	1.34	1.69	1.48
Fruits	22.40	15.78	29.98	2.68	1.03	2.28
Branches/trunk	20.70	28.38	26.44	4.76	3.80	4.22
Roots	11.60	20.82	20.20	2.85	2.98	2.96
Total	78.70	114.78	113.72	11.64	9.52	10.95

High N accumulation with ACPS than CMP but P accumulation similar for all practices.


ACPS Nutrition Studies (3)



Leaf NPK concentration (%) determined in June 2009 at Immokalee.

- Sufficient NPK concentrations.
- Drip OHS was effective in enhancing N uptake compared with the other two irrigation methods studied.
- Leaf P concentration was high (0.17-0.30%) in all treatments
- Leaf K concentration was within optimum and high ranges (1.2-2.4%) suggesting no significant differences between ACPS and Conventional method.


ACPS Nutrition Studies (4)



- Greater inorganic N in irrigated than non-irrigated zones
- Better N contents in irrigated zones of ACPS than Conventional.

Ammonium and nitrate distribution in the irrigated and non-irrigated zone


ACPS Nutrition Studies (5)



- Greater P in irrigated than non-irrigated zones
- Soil P contents in irrigated zones of Drip greater than Conventional.

Soil P distribution in the irrigated and non-irrigated zones



ACPS Nutrition Studies (6)



**Higher NPK conc.
in irrigated vs.
non-irrigated zones
of drip fertigation.
D=area below
dripper, T=tree**

**Lateral ammonium-N (mg/kg) distribution in July 2010
at the Lake Alfred site using drip fertigation.**


ACPS Nutrition Studies (7)



Vertical P distribution at Immokalee and Lake Alfred sites in 2010

Less P leaching with OHS than CMP in 2010.
High P at Lake Alfred than Immokalee

ACPS Nutrition Studies (8)




**ACPS fertigation
had greater tree size
than conventional
practice**

**Canopy volume as a function of fertilization
practice at the Lake Alfred site**


ACPS Nutrition Studies (9)

**Lateral RLD (cm cm^{-3})
distribution using CMP**



T=tree, M=microsprinkler
Roots uniformly distributed
around the tree

**Lateral RLD (cm cm^{-3})
distribution using DOHS**




T=tree, D=dripper, Roots
concentrated below the
drippers

ACPS Nutrition Studies (7)

Positions in the irrigated zones of showed higher root density than non-irrigated zones

M=microsprinkler

T=tree



Lateral root density (cm cm⁻³) distribution using ACPS microsprinkler

Summary

- Daily, frequent irrigation critical for improved tree performance, soil moisture distribution and water use.
- HLB affected trees use 22 to 35% less water than the non-affected trees.
- ACPS practices could be adapted to grower practices for vigorous tree growth, water use, greater root density and nutrient accumulation.

Acknowledgements

- Dr. Morgan, Dr. Schumann, Dr. Ebel, Dr. Hamido – University of Florida
- Grove/Orchard Space:
 - UF/IFAS SWFREC, Immokalee, FL
 - Gapway Groves, Auburndale, FL
 - Pacific Inc., Ave Maria, FL
 - Orange Co, Arcadia, FL
 - Ben Hill Griffin, Avon Park, FL
- Funding: Southwest FL WMD, FDACS, UF/IFAS
- **Thanks to Mrs. Ana Villalpando and Dr. Medina for the invitation.
Muchas Gracias!**

Muchas gracias!!